

SELECTED SOURCES OF INFORMATION

Working Group on Articles 9 and 10 of the WHO FCTC

(April 2012)

1) Product Characteristics in relation to Fire-Risk (Reduced Ignition Propensity)

General

Mueller J (2006). Cigarette fire safety standards, figure 2.4. In: *Second international conference on fire safer cigarettes*. Boston, Massachusetts, Harvard School of Public Health.

WHO Study Group on Tobacco Product Regulation (2008). The Scientific Basis of Tobacco Product Regulation. Second report of a WHO Study Group. Advisory note on 'fire-safer' cigarettes: approaches to reduced ignition propensity. *WHO Technical Report Series*, no. 951. Available at:

www.who.int/entity/tobacco/global_interaction/tobreg/publications/9789241209519.pdf

Regulatory Requirements and Standards

Australian Government ComLaw (2008). Trade Practices (Consumer Product Safety Standard) (Reduced Fire Risk Cigarettes) Regulations 2008. Select Legislative Instrument 2008 No. 195. *Federal Register of Legislative Instruments F2008L03475*. Available at: www.comlaw.gov.au/Details/F2008L03475/Download

Canada (2005). Tobacco Act, Cigarette Ignition Propensity Regulations P.C. 2005-1125, June 7, 2005. *Canada Gazette*, Part II, Vol. 139, No. 13, p. 1505. Available at: www.gazette.gc.ca/archives/p2/2005/2005-06-29/pdf/g2-13913.pdf

Commission Implementing Decision 2011/496/EU on the compliance of standard EN 16156:2010 'Cigarettes — Assessment of the ignition propensity — Safety requirement' and of standard EN ISO 12863:2010 'Standard test method for assessing the ignition propensity of cigarettes' with the general safety requirement of Directive 2001/95/EC of the European Parliament and of the Council and publication of the references of standard EN 16156:2010 'Cigarettes — Assessment of the ignition propensity — Safety requirement' and of standard EN ISO 12863:2010 'Standard test method for assessing the ignition propensity of cigarettes' in the Official Journal of the European Union. *Official Journal of the European Union*, L 205, 10.8.2011, p. 31. Available at: <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2011:205:0031:0032:EN:PDF>

European Committee for Standardization (CEN) (2010). Cigarettes – Assessment of the ignition propensity – Safety requirement. CEN standard EN 16156:2010.

Government of South Africa, Department of Health (1993). Notice on the tobacco products control act (Act No. 83 of 1993)- Regulations relating to the standards for manufacturing of reduced ignition propensity (RIP) cigarettes. Government Notice N° R.429, 16 May 2011. *Staatskoerant*, 16 Mei 2011, Gazette No. 34302, p. 3. Available at: www.tobaccocontrollaws.org/files/live/South%20Africa/South%20Africa%20-%20RIP%20Regs%20-%20national.pdf

New York State. Fire Safety Standards for Cigarettes, Part 429, Chapter XII, Title 19 of the New York Code of Rules and Regulations (NYCRR). Available at: <http://government.westlaw.com/linkedslice/default.asp?SP=nycrr-1000>

Test methods

ASTM International (2009). Standard Test Method for Measuring the Ignition Strength of Cigarettes. ASTM International Standard E2187-09.

International Organization for Standardization (2010). Standard test method for assessing the ignition propensity of cigarettes. ISO 12863:2010.

International Organization for Standardization (2010). Standard test method for assessing the ignition propensity of cigarettes. EN ISO 12863:2010

Standards Australia (2007). Determination of the extinction propensity of cigarettes. AS 4830-2007.

Standards New Zealand (2009). Determination of the extinction propensity of cigarettes. NZS/AS 4830-2007.

Effects of cigarette fires on health and lives, and fire reduction by RIP cigarettes

Alpert HR, O'Connor RJ, Spalletta R, Connolly GN (2010). Recent advances in cigarette ignition propensity research and development. *Fire Technology*, 46(2):275-289. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2873202/pdf/nihms122218.pdf>

Australian Competition & Consumer Commission (2008). Regulation impact statement - the regulation of reduced fire risk cigarettes under the trade practices act 1974. Available at:

www.accc.gov.au/content/item.phtml?itemId=807543&nodeId=aac0bdf39e1ba1d76a42e81eedf77ab1&fn=Reduced+fire+risk+cigarettes%E2%80%94regulation+impact+statement%E2%80%94July+2008.doc

Australian Government ComLaw (2008). Trade Practices (Consumer Product Safety Standard) (Reduced Fire Risk Cigarettes) Regulations 2008, No. 195 as amended. Available at: www.comlaw.gov.au/Details/F2009C00252

Botkin JR (1988). The fire-safe cigarette. *Journal of the American Medical Association*, 260(2):226-9.

Brunnemann KD, Hoffmann D, Gairola CG, Lee BC (1994). Low ignition propensity cigarettes: smoke analysis for carcinogens and testing for mutagenic activity of the smoke particulate matter. *Food Chemical Toxicology*, 32(10):917-922. Available at: http://tobaccodocuments.org/product_design/2029138624-8638.html

Canada (2005). Cigarette Ignition Propensity Regulations. Tobacco Act. SOR/2005-178. <http://laws-lois.justice.gc.ca/PDF/SOR-2005-178.pdf>

Chapman S, Balmain A (2004). Reduced-Ignition Propensity Cigarettes: A review of policy relevant information. Report prepared for the (Australia) Commonwealth Department of Health and Ageing. Available at: [http://www.health.gov.au/internet/main/publishing.nsf/Content/1526AD854C903649CA256F8E0011ACBE/\\$File/smoking_rip.pdf](http://www.health.gov.au/internet/main/publishing.nsf/Content/1526AD854C903649CA256F8E0011ACBE/$File/smoking_rip.pdf)

Connolly GN, Alpert HR, Rees V, Carpenter C, Wayne GF, Vallone D, Koh H (2005). Effect of the New York State cigarette fire safety standard on ignition propensity, smoke constituents, and the consumer market. *Tobacco Control*, 14:312-327. Available at : <http://tobaccocontrol.bmj.com/content/14/5/321.full.pdf+html>

Côté F, Létourneau C, Mullard G, Voisine R (2010). Estimation of nicotine and tar yields from human-smoked cigarettes before and after the implementation of the cigarette ignition propensity regulations in Canada. *Regulatory Toxicology and Pharmacology*, 2011;61(3 Suppl):S51–59. Available at: http://pdn.sciencedirect.com/science?_ob=ImageURL&_cid=272321&_user=8658645&_pii=S0273230010000449&_check=y&_origin=article&_zone=toolbar&_coverDate=01-Dec-2011&view=c&originContentFamily=serial&wchp=dGLbVIB-zSkWz&md5=92931353de24adfaa2b29eeca33d8eb/1-s2.0-S0273230010000449-main.pdf

European Commission (2004). Tobacco or Health in the European Union. Past, present and future. The ASPECT Consortium. Available at: http://ec.europa.eu/health/ph_determinants/life_style/Tobacco/Documents/tobacco_exs_en.pdf

Gann RG (2007). Measuring the ignition propensity of cigarettes. *International Interflam Conference*, 11th Proceedings. Volume 1. September 3-5, 2007, London, England, 145-155. Available at: www.fire.nist.gov/bfrlpubs/fire07/PDF/f07068.pdf

Gormley Michael (2005). 1st year of fire safe cigarette law yields fewer deaths. *Associated Press*. Available at: <http://ash.org/no-smoking/nov05/11-22-05-2.html>

Gunja M, Ferris Wayne G, Landman A, Connolly G, McGuire A (2002). The case for fire safe cigarettes made through industry documents. *Tobacco Control*, 11:346-353. Available at: <http://tobaccocontrol.bmj.com/content/11/4/346.full.pdf>

Health Canada (2004). Economic evaluation of health canada's regulatory proposal for reducing fire risks from cigarettes. Available at: www.hc-sc.gc.ca/hc-ps/pubs/tobac-tabac/evaluation-risks-risques/index-eng.php

Information letter of 19 May 2011 from the Finnish Ministry of the Interior to the European Commission

Laugesen M, Duncanson M, Fraser T, McClellan V, Linehan B, Shirley R (2003). Hand rolling cigarette papers as the reference point for regulating cigarette fire safety. *Tobacco Control*, 12(4):406-410. Available at: www.ncbi.nlm.nih.gov/pmc/articles/PMC1747792/pdf/v012p00406.pdf

O'Connor RJ, Fix BV, Hammond D, Giovino GA, Hyland A, Fong GT, Cummings KM (2010). The impact of reduced ignition propensity cigarette regulation on smoking behaviour in a cohort of Ontario smokers. *Injury Prevention*, 16(6):420-422.

O'Connor RJ, Rees VW, Norton KJ, Cummings KM, Connolly GN, Alpert HR, Sjödin A, Romanoff L, Li Z, June KM, Giovino GA (2010). Does switching to reduced ignition propensity cigarettes alter smoking behavior or exposure to tobacco smoke constituents? *Nicotine & Tobacco Research*, 12(10):1011-1018.

O'Connor RJ, Bauer JE, Giovino GA, Hammond D, Hyland A, Fong GT, Cummings KM. (2007). Prevalence of behaviors related to cigarette-caused fires: a survey of Ontario smokers. *Injury Prevention*, 13(4):237-242.

O'Connor RJ, Giovino GA, Fix BV, Hyland A, Hammond D, Fong GT, Bauer U, Cummings KM. (2006). Smokers' reactions to reduced ignition propensity cigarettes. *Tobacco Control*, 15(1):45-49.

Office of the Deputy Prime Minister; London (UK) (report prepared by BTTG Ltd) (2004). A programme of work to compare the propensity of fire safe cigarettes and conventional cigarettes to ignite textile materials used in a domestic environment.

Available at: <http://tobacco.health.usyd.edu.au/assets/pdfs/Ingredients-and-Regulations/ODPM-PROJECT-DRAFT-REPORT-3rd-revision.pdf>

Palmiere C, Staub C, La Harpe R, Mangin P (2009). Ignition of a human body by a modest external source: a case report. *Forensic Science International*, 188(1-3):e17-19.

Patskan G et al (2000). Toxicological characterization of a novel cigarette paper. Society of Toxicology Annual Meeting 2000. Available at:

<http://legacy.library.ucsf.edu/tid/krt95c00/pdf;jsessionid=AF5CD5A5838EFE4F726BCB91023B4E79.tobacco03>

Rustemeier K, Stabbert R, Haussmann HJ, Roemer E, Carmines EL (2002). Evaluation of the potential effects of ingredients added to cigarettes. Part 2: chemical composition of mainstream smoke. *Food Chemical Toxicology*, 40(1):93-104.

Seidenberg AB, Rees VW, Alpert HR, O'Connor RJ, Connolly GN (2011). Ignition strength of 25 international cigarette brands. *Tobacco Control*, (1):77-80.

Tohamy SM, Aiken DV (2007). Assessing regulatory effectiveness with exogenously declining risk: a case study of the CPSC's 1973 mattress standard. *Journal of Safety Research*, 8(6):661-668.

WHO Study Group on Tobacco Product Regulation (2008). The Scientific Basis of Tobacco Product Regulation. Second report of a WHO Study Group. Advisory note on 'fire-safer' cigarettes: approaches to reduced ignition propensity. *WHO Technical Report Series*, no. 951. Available at:

www.who.int/entity/tobacco/global_interaction/tobreg/publications/9789241209519.pdf

2) Public Disclosure – Toxic Constituents and Emissions

Decision FCTC/COP4(7). Guidelines for the implementation of Article 12 of the WHO Framework Convention on Tobacco Control (Education, communication, training and public awareness). In: Decisions and ancillary. Fourth session of the Conference of the Parties to the WHO Framework Convention on Tobacco Control, Punta del Este, Uruguay, 15–20 November 2010. FCTC/COP/4/REC/1, p. 8-26. Available at: http://apps.who.int/gb/fctc/PDF/cop4/FCTC_COP4_REC1.pdf

Decision FCTC/COP3(10). Guidelines for implementation of Article 11 (Packaging and labelling of tobacco products). In: Decisions and ancillary. Third session of the Conference of the Parties to the WHO Framework Convention on Tobacco Control, Durban, South Africa, 17–22 November 2008. FCTC/COP/3/REC/1, p. 16-28. Available at: http://apps.who.int/gb/fctc/PDF/cop3/FCTC_COP3_REC1-en.pdf

European Commission, Health and Consumer Protection Directorate-General (2007). Reporting on tobacco product ingredients: practical guide (Brussels, 31 May 2007) (SANCO C6 TPE/ub D(2007) 360206). Available at: http://ec.europa.eu/health/ph_determinants/life_style/Tobacco/Documents/practical_guidance_en.pdf

European Parliament and the Council of the European Union (2001). Directive 2001/37/EC of the European Parliament and of the Council of 5 June 2001 on the approximation of the laws, regulations and administrative provisions of the Member States concerning the manufacture, presentation and sale of tobacco products. Available at: <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2001:194:0026:0034:EN:PDF>

Government of Belgium. Search tool for tobacco product information (last consulted April, 2012) Available at: www.health.belgium.be/pls/apex/f?p=EMTOC_SEARCH:1:115199926172735

International Agency for Research on Cancer (IARC) (2008). *Handbooks of Cancer Prevention, Volume 12, Part 5.3 – Methods for Evaluating Tobacco Control Policies: Measures to Assess the Effectiveness of Tobacco Product Regulation*. World Health Organization, Geneva, 2008. Available at: www.iarc.fr/en/publications/pdfs-online/prev/handbook12/Tobacco_vol12_5C.pdf

Ipsos-Eureka (2009). Public health value of disclosed cigarette ingredients and emissions data (prepared for the Department of Health and Ageing, Australia, January 2009). Available at: www.health.gov.au/internet/main/publishing.nsf/Content/phd-tobacco-phv-cig-ing-em-data

McNeill A, Joossens L, Jarvis M (commissioned by Action on Smoking and Health London) (2004). Review of the implementation of the Tobacco Product Regulation Directive 2001/37/EC. Available at: www.ash.org.uk/files/documents/ASH_164.pdf

Report by WHO's Tobacco Free Initiative. Work in progress in relation to Articles 9 and 10 of the WHO Framework Convention on Tobacco Control (2010). Conference of the Parties to the WHO Framework Convention on Tobacco Control, fourth session, Punta del Este, Uruguay, 15-20 November 2010, FCTC/COP/4/INF.DOC./2 (15 September 2010). Available at: (http://apps.who.int/gb/fctc/PDF/cop4/FCTC_COP4_ID2-en.pdf

3) Background Paper on Tobacco Addictiveness Reduction

Paragraph 2

Royal College of Physicians (2000). Nicotine addiction in Britain: a report of the tobacco advisory group of the Royal College of Physicians. *Royal College of Physicians of London*. Available at: <http://bookshop.rcplondon.ac.uk/contents/pub131-2740272e-57b4-4567-b87c-cda5ef7cd7e0.pdf>

Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR) (2010). Addictiveness and attractiveness of tobacco additives. European Commission DG Health & Consumers. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenihr_o_031.pdf

World Health Organization. Lexicon of alcohol and drug terms published by the World Health Organization.
www.who.int/substance_abuse/terminology/who_lexicon/en/

WHO Study Group on Tobacco Product Regulation (2007). The Scientific Basis of Tobacco Product Regulation: Report of a WHO study group, *WHO Technical Report Series*, no. 945. Available at: http://www.who.int/tobacco/global_interaction/tobreg/9789241209458.pdf

WHO Study Group on Tobacco Product Regulation (2012). Report on the Scientific Basis of Tobacco Product Regulation: Fourth Report of a WHO Study Group, *WHO Technical Report Series*, no. 967. Available at: http://whqlibdoc.who.int/publications/2012/9789241209670_eng.pdf

Paragraph 3

Belluzzi JD, Wang R, Leslie FM (2005). Acetaldehyde enhances acquisition of nicotine self-administration in adolescent rats. *Neuropsychopharmacology*, 30:705-712.

Berggren U, Eriksson M, Fahlke C, Blennow K, and Balldin J (2007). Different effects of smoking or use of smokeless tobacco on platelet MAO-B activity in type 1 alcohol-dependent subjects. *Alcohol and Alcoholism*, 42(3):267 - 271. Available at: <http://alcalc.oxfordjournals.org/content/42/3/267.full.pdf+html>

Clemens KJ, Caillé, S, Stinus, L, Cador M (2009). The addition of five minor tobacco alkaloids increases nicotine-induced hyperactivity, sensitization and intravenous self-administration in rats. *International Journal of Neuropsychopharmacology*, 15:1-12.

Dixon M, Lambing K, Seeman J (2000). On the Transfer of Nicotine from Tobacco to the Smoker. A brief review of ammonia and "pH" factors. *Beitraege zur Tabakforschung / Contributions to Tobacco Research*, 19:103-113.

Fowler JS, Volkow ND, Wang GJ, Pappas N, Logan J, MacGregor R, Alexoff D, Shea C, Schlyer D, Wolf AP, Warner D, Zezulkova I, Cilento R (1996). Inhibition of monoamine oxidase B in the brains of smokers. *Nature*, 379:733-736.

Fowler JS, Volkow ND, Wang GJ, Pappas N, Logan J, Shea C, Alexoff D, MacGregor RR, Schlyer DJ, Zezulkova I, Wolf AP (1996). Brain monoamine oxidase inhibition in cigarette smokers. *Proceedings of the National Academy of Sciences of the United States of America*, 93:14065-14069. Available at: www.pnas.org/content/93/24/14065.full.pdf+html

Fowles J (2001). Chemical Factors Influencing the Addictiveness and Attractiveness of Cigarettes in New Zealand - Final Report, prepared as Part of a New Zealand Ministry of Health Contract for Scientific Services. Available at: [www.moh.govt.nz/moh.nsf/pagescm/1004/\\$File/chemicalfactorsaddictivenesscigarettes.pdf](http://www.moh.govt.nz/moh.nsf/pagescm/1004/$File/chemicalfactorsaddictivenesscigarettes.pdf)

Fowles J, Bates M (2000). The chemical constituents of cigarettes and cigarette smoke: Priorities for harm reduction. A report by the Institute of Environmental Science and Research for the New Zealand Ministry of Health. Available at: [www.moh.govt.nz/moh.nsf/pagescm/1003/\\$File/chemicalconstituentscigarettespriorities.pdf](http://www.moh.govt.nz/moh.nsf/pagescm/1003/$File/chemicalconstituentscigarettespriorities.pdf)

Henningfield JE, Pankow J, Garrett B (2004). Ammonia and other chemical base tobacco additives and cigarette nicotine delivery: issues and research needs. *Nicotine & Tobacco Research*, 6:199-205.

Herraiz T, Chaparro C (2005). Human monoamine oxidase is inhibited by tobacco smoke: β -carboline alkaloids act as potent and reversible inhibitors. *Biochemical and Biophysical Research Communications*, 326(2):378-386

Hughes JR, Rose GL, Callas PW (2000). Nicotine is more reinforcing in smokers with a past history of alcoholism than in smokers without this history. *Alcoholism: Clinical and experimental Research*, 24(11):1633-1638.

Khalil AA, Steyn S, Castagnoli N Jr (2000). Isolation and characterization of a monoamine oxidase inhibitor from tobacco leaves, *Chemical Research in Toxicology*, 13, 31–35. Available at: <http://ipac.kacst.edu.sa/edoc/eBook/1911.pdf>

Talhout R, Opperhuizen A, Van Amsterdam JGC (2007). Role of acetaldehyde in tobacco smoke addiction. *European Neuropsychopharmacology* 17(10):627-636.

Van Amsterdam J, Sleijffers A, van Spiegel P, Blom R, Witte M, van de Kasstele J, Blokland M, Steerenberg P, Opperhuizen A (2011). Effect of ammonia in cigarette tobacco on nicotine absorption in human smokers. *Food Chemical Toxicology*, 49(12):3025-3030.

Wrona MZ, Wakiewicz J, Han QP, Han J, Li H, Dryhurst G (1997). Putative oxidative metabolites of 1-methyl-6-hydroxy-1,2,3,4-tetrahydro-beta-carboline of potential relevance to the addictive and neurogenerative consequences of ethanol abuse. *Alcohol*, 14(3):213-223.

Paragraph 4

Benowitz NL (1989). Dosimetric studies of compensatory cigarette smoking; In: *Independent Scientific Committee on Smoking and Health: Nicotine, Smoking, and the Low Tar Programme*; London, November 18- 20, 1986, N. Wald and P. Froggatt (Editors). Oxford: Oxford University Press, pp. 133-150.

Benowitz NL, Hall SM, Herning RI, Jacob P 3rd, Jones RT, Osman AL (1983). Smokers of low-yield cigarettes do not consume less nicotine. *New England Journal of Medicine*, 309:139-142.

Hatsukami DK, Le CT, Zhang Y, Joseph AM, Mooney ME, Carmella SG, Hecht SS (2006). Toxicant exposure in cigarette reducers versus light smokers. *Cancer Epidemiology, Biomarkers & Prevention*, Dec;15(12):2355-2358. Available at: <http://cebp.aacrjournals.org/content/15/12/2355.full.pdf>

Hatsukami DK, Perkins KA, Lesage MG, Ashley DL, Henningfield JE, Benowitz NL, Backinger CL, Zeller M (2010). Nicotine reduction revisited: science and future directions. *Tobacco Control*, 19(5):e1-e10.

Kozlowski LT, Mehta NY, Sweeney CT, Swartz SS, Vogler GP, Jarvis M.J., Wes, RJ (1998). Filter ventilation and nicotine content of tobacco in cigarettes from Canada and the United States. *Tobacco Control*, 7:369-375. Available at: <http://tobaccocontrol.bmj.com/content/7/4/369.full.pdf>

O'Connor RJ, Cummings KM, Giovino GA, McNeill A, Kozlowski LT (2006). How did UK cigarette makers reduce tar to 10 mg or less? *British Medical Journal*, 332 (7536):302. Available at: <http://toxicology.usu.edu/endnote/302.pdf>

Paragraph 5

American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4th ed. Washington, DC, 1994.

Carter LP, Stitzer ML, Henningfield JE, O'Connor RJ, Cummings KM, Hatsukami DK (2009). Abuse liability assessment of tobacco products including potential reduced exposure products. *Cancer Epidemiology, Biomarkers & Prevention*, 18(12):3241-3262. Available at: <http://cebp.aacrjournals.org/content/18/12/3241.full.pdf+html>

Donny EC, Dierker LC (2007). The absence of DSM-IV nicotine dependence in moderate-to-heavy daily smokers. *Drug and Alcohol Dependence*, 89(1):93-96. Available at: www.ncbi.nlm.nih.gov/pmc/articles/PMC1924493/pdf/nihms22795.pdf

Fagerstrom KO (2011). Determinants of Tobacco Use and Renaming the FTND to the Fagerstrom Test for Cigarette Dependence. *Nicotine and Tobacco Research*, 14(1):75-8.

Health Canada. (2007). Guidance Document - Clinical Assessment of Abuse Liability for Drugs with Central Nervous System Activity. Available at: www.hc-sc.gc.ca/dhp-mps/alt_formats/hpfb-dgpsa/pdf/prodpharma/abuse_liabilityabusifusageclin-eng.pdf

Heatherton TF, Kozlowski LT, Frecker RC, Fagerstrom KO (1991). The Fagerstrom Test for Nicotine Dependence: A revision of the Fagerstrom Tolerance Questionnaire. *British Journal of Addiction*, 86:1119-1127.

Hughes JR (2006). Should criteria for drug dependence differ across drugs? *Addiction*, 101 (Suppl. 1):134-141.

Paragraph 6

Cooperation Centre for Scientific Research Relative to Tobacco (CORESTA). CORESTA Recommended Methods No. 35 (August 2010). Determination of total alkaloids (as nicotine) in tobacco by continuous flow analysis. Available at: [www.coresta.org/Recommended_Methods/CRM_35-update\(Aug10\).pdf](http://www.coresta.org/Recommended_Methods/CRM_35-update(Aug10).pdf) and No. 62 (February 2005). Determination of nicotine in tobacco and tobacco products by gas chromatographic analysis. Available at: www.coresta.org/Recommended_Methods/CRM_62.pdf

Health Canada (1999). Official Method T-301, Determination of Alkaloids in Whole Tobacco, December 1999. Available at: www.hc-sc.gc.ca/hc-ps/alt_formats/hecs-sesc/pdf/tobac-tabac/legislation/reg/indust/method/whole-entier/alkaloids-eng.pdf

Severson RF, McDuffie KL, Arrendale RF, Gwynn GR, Chaplin JF, Johnson AW (1981). Rapid method for the analysis of tobacco nicotine alkaloids. *Journal of Chromatography*, 211:111-121.

Stanfill SB, Jia LT, Watson CH (2009). Rapid and chemically-selective quantification of nicotine in smokeless tobacco products using gas chromatography/mass spectrometry. *Journal of Chromatographic Science*, 47:902-909.

Paragraph 7

Ayo-Yusuf OA, Swart TJP, Pickworth WB (2004). Nicotine delivery capabilities of smokeless tobacco products and implications for control of tobacco dependence in South Africa. *Tobacco Control*, 13:186–189. Available at:

<http://tobaccocontrol.bmj.com/content/13/2/186.full.pdf+html>

Djordjevic MV, Doran KA (2009). Nicotine Content and Delivery Across Tobacco Products, In: J.E. Henningfield et al. (eds.), *Nicotine Psychopharmacology, Handbook of Experimental Pharmacology* 192, National Health Institute, pp. 61-82.

Fant RV, Henningfield JE, Nelson RA (1999). Pharmacokinetics and pharmacodynamics of moist snuff in humans. *Tobacco Control*, 8:387–392. Available at:

<http://tobaccocontrol.bmj.com/content/8/4/387.full.pdf+html>

Hadidi KA, Mohammed FI (2004). Nicotine content in tobacco used in hubble-bubble smoking. *Saudi Medical Journal*, 25 (7):912-917.

Idris AM, Nair, J, Oshima H, Friesen M, Bronet I, Faustman EM, Bartsch H (1991). Unusually high levels of carcinogenic tobacco-specific nitrosamines in Sudan snuff (*toombak*). *Carcinogenesis*, 12:1115–1118.

Incekara F (1979). *Industrial Plants, Vol. 4 (Pleasurable Plants)*. Aegean University Faculty of Agriculture Press. No:84 Bornova, İzmir, Turkey.

International Agency for Research on Cancer (IARC)(2004). IARC Monographs on the Evaluation of Carcinogenic Risks to Humans Volume 83: Tobacco Smoke and Involuntary Smoking, and volume 85: Betel-quid and Areca-nut Chewing and Some Areca-nut-derived Nitrosamines. Available at:

<http://monographs.iarc.fr/ENG/Monographs/vol83/volume83.pdf>

<http://monographs.iarc.fr/ENG/Monographs/vol85/index.php>

Malson JL, Sims K, Murty R, Pickworth WB (2001). Comparison of the nicotine content of tobacco used in bidis and conventional cigarettes. *Tobacco Control*, 10:181–183.

Available at: <http://tobaccocontrol.bmj.com/content/10/2/181.full.pdf>

Perfetti TA, Norman AB, Gordon BM, Coleman WM III, Morgan WT, Dull GM, Miller CW (2000). The transfer of nicotine from nicotine salts to mainstream smoke. *Beitraege zur Tabakforschung / Contributions to Tobacco Research*, 19(3):141-160.

Stanfill SB, Connolly GN, Zhang L, Jia LT, Henningfield JE, Richter P, Lawler TS, Ayo-Yusuf OA, Ashley DL, Watson CH (2011). Global surveillance of oral tobacco products: total nicotine, unionised nicotine and tobacco-specific N-nitrosamines. *Tobacco Control*, 20(3):e2.

Tso TC (1990). Organic metabolism- alkaloids. In: Tso TC (ed) *Production, physiology and biochemistry of tobacco plant*. IDEALS, Beltsville, MD, pp. 427–486.

Paragraph 8

Benowitz NL (2008). Clinical pharmacology of nicotine: Implications for understanding, preventing, and treating tobacco addiction. *Clinical Pharmacology & Therapeutics*, 83(4):531–541.

Brown & Williamson (1989). Study of the tobacco blend composition of Next and Merit de-nic cigarettes/151. Bates No. 599005141-5150. Available at: <http://legacy.library.ucsf.edu/tid/gxi41f00/pdf>

Brown & Williamson (1989). Evaluations of PM's Nicotine Free Cigarettes, Bates No. 599005151-5155. Available at: <http://legacy.library.ucsf.edu/tid/hxi41f00/pdf>

Chen J, Higby R, Tian D, Tan D, Johnson MD, Xiao Y, Kellar KJ, Feng S, Shields PG (2008). Toxicological analysis of low-nicotine and nicotine-free cigarettes. *Toxicology*, 249(2-3):194-203.

Djordjevic MV, Sigountos CW, Brunnemann KD Hoffmann D (1990). Tobacco-specific nitrosamine delivery in the mainstream smoke of high- and low-yield cigarettes smoked with varying puff volume. In: *CORESTA Symposium Information Bulletin*. Hellas, October 7–11, 1990, p. 209.

Djordjevic MV (2010). Nicotine content in tobacco and tobacco smoke. In: Boyle P, Gray N, Henningfield J, Seffrin J, Zatonski W. *Tobacco: Science, policy and public health*. Oxford University Press, Oxford, pp. 197-220.

Dunsby J, Bero L (2004). A nicotine delivery device without the nicotine? Tobacco industry development of low nicotine cigarettes. *Tobacco Control*, 13:362-369
Hansen J, Muller J (1973). Report on no-nicotine project. R.J. Reynolds Collection. Bates No. 509386784-6801. Available at: <http://legacy.library.ucsf.edu/tid/gbn73d00/pdf>

Philip Morris USA (1994). Alkaloid Reduced Tobacco (ART) Program. Bates No. 2057908259-8291. Available at: <http://legacy.library.ucsf.edu/tid/top46b00/pdf>

Philip Morris (1987). Process and apparatus for the semicontinuous extraction of nicotine from tobacco. Bates No. 2028585410-5445. Available at: <http://legacy.library.ucsf.edu/tid/zfs04e00/pdf>

Philip Morris (1999). Philip Morris developments: The denicotinized tobacco story. Bates No. 2078425674-5792. Available at: <http://legacy.library.ucsf.edu/tid/cbp75c00/pdf>

R.J. Reynolds (1983) Comparison of genetically low nicotine tobaccos with denicotinized tobaccos. Bates No. 501660695-0704. Available at:
<http://legacy.library.ucsf.edu/tid/xbl39d00/pdf>

Strasser AA, Lerman C, Sanborn PM, Pickworth WB, Feldman EA (2007). New lower nicotine cigarettes can produce compensatory smoking and increased carbon monoxide exposure. *Drug and Alcohol Dependence*, 86(2-3):294-300.

Valleau WD (1949). Breeding low-nicotine tobacco. *Journal of Agricultural Research*, 78:171–181.

Wang P, Liang Z, Zeng J, Li W, Sun X, Miao Z and Tang K (2008). Generation of tobacco lines with widely different reduction in nicotine levels via RNA silencing approaches. *Journal of Biosciences*, 33:177–184. Available at:
www.ias.ac.in/jbiosci/jun2008/177.pdf

Xie J, Song W, Maksymowicz W, Jin W, Cheah K, Chen W, Carnes C, Ke J, Conkling MA (2004). Biotechnology: A Tool for Reduced Risk Tobacco Products – The Nicotine Experience from Test Tube to Cigarette Pack. *Recent Advances in Tobacco Science*, 30:17-37.

Paragraph 9

Hatsukami DK, Perkins KA, Lesage MG, Ashley DL, Henningfield JE, Benowitz NL, Backinger CL, Zeller M (2010). Nicotine reduction revisited: science and future directions. *Tobacco Control*, 19(5):e1-e10.

Henningfield JE, Benowitz NL, Slade J, Houston TP, Davis RM, Deitchman SD (1998). Reducing the addictiveness of cigarettes. Council on Scientific Affairs, American Medical Association. *Tobacco Control*, 7(3):281-293. Available at:
<http://tobaccocontrol.bmj.com/content/7/3/281.full.pdf+html>

Paragraph 10

Benowitz NL, Dains KM, Hall SM, Stewart S, Wilson M, Dempsey D, Jacob P III (2012). Smoking behavior and exposure to tobacco toxicants during 6 months of smoking progressively reduced nicotine content cigarettes. *Cancer Epidemiology, Biomarkers & Prevention*. (Epub ahead of print, Mar 15).

Benowitz NL, Hall SM, Stewart S, Wilson M, Dempsey D, Jacob P III (2007). Nicotine and carcinogen exposure with smoking of progressively reduced nicotine content cigarette. *Cancer Epidemiology, Biomarkers & Prevention*, 16(11):2479–2485. Available at: <http://cebp.aacrjournals.org/content/16/11/2479.full.pdf>

Hammond D, Fong GT, Cummings KM, Hyland A (2005). Smoking topography, brand switching, and nicotine delivery: Results from an *in vivo* Study. *Cancer Epidemiology, Biomarkers & Prevention*, 14(6):1370-1375. Available at: <http://cebp.aacrjournals.org/content/14/6/1370.full.pdf+html>

Rose JE, Behm FM (2004). Effects of low nicotine content cigarettes on smoke intake. *Nicotine & Tobacco Research*, 6(2):309-319.

Paragraph 11

Hatsukami DK, Perkins KA, Lesage MG, Ashley DL, Henningfield JE, Benowitz NL, Backinger CL, Zeller M (2010). Nicotine reduction revisited: science and future directions. *Tobacco Control*, 19(5):e1-e10.

Hatsukami DK, Kotlyar M, Hertsgaard LA, Zhang Y, Carmella SG, Jensen JA, Allen SS, Shields PG, Murphy SE, Stepanov I, Hecht SS (2010). Reduced nicotine content cigarettes: effects on toxicant exposure, dependence and cessation. *Addiction*, 105(2):343-355.

Henningfield JE, Benowitz NL, Connolly GN, Davis RM, Gray N, Myers ML, Zeller M (2004). Reducing tobacco addiction through tobacco product regulation. *Tobacco Control*, 13:132-135.

Henningfield JE, Benowitz NL, Slade J, Houston TP, Davis RM, Deitchman SD (1998). Reducing the addictiveness of cigarettes. Council on Scientific Affairs, American Medical Association. *Tobacco Control*, 7(3):281-293.

Henningfield, JE, Zeller, M (2002). Could science-based regulation make tobacco products less addictive? *Yale Journal of Health Policy, Law and Ethics*, III:127–138.

Paragraph 12

Hatsukami DK, Biener L, Leischow SJ, Zeller MR (2012). Tobacco and Nicotine Product Testing. *Nicotine & Tobacco Research*, 14(1):7-17.

Hatsukami DK, Perkins KA, Lesage MG, Ashley DL, Henningfield JE, Benowitz NL, Backinger CL, Zeller M (2010). Nicotine reduction revisited: science and future directions. *Tobacco Control*, 19(5):e1-e10.

Hecht SS (2012). Research opportunities related to establishing standards for tobacco products under the family smoking prevention and tobacco control act. *Nicotine & Tobacco Research*, 14(1):18-28.

Hecht SS, Yuan J-M, Hatsukami DK (2010). Applying tobacco carcinogen and toxicant biomarkers in product regulation and cancer prevention. *Chemical Research in Toxicology*, 23:1001–1008. Available at:
www.ncbi.nlm.nih.gov/pmc/articles/PMC2891118/pdf/nihms198929.pdf

Tengs TO, Ahmad S, Savage JM, Moore R, Gage E (2005). The AMA proposal to mandate nicotine reduction in cigarettes: a simulation of the population health impacts. *Preventive Medicine*, 40:170–180. Available at:
http://unlv.academia.edu/SajjadAhmad/Papers/972756/The_AMA_proposal_to_mandate_nicotine_reduction_in_cigarettes_a_simulation_of_the_population_health_impacts